

NEWS & VIEWS

Open Access

Spatiotemporal pulse weaving scalar optical hopfions

Chenhui Li¹, Sicong Wang¹ and Xiangping Li¹✉

Abstract

Scalar optical hopfions weaved by nested equiphase lines in the shape of a toroidal vortex are theoretically designed and experimentally demonstrated. This category of hopfions manifesting as a spatiotemporally structured pulse propagating in space-time may enable encoding and transferring optical topological information in an additional (temporal) dimension.

Topological solitons with topologically protected spin textures are cutting-edge researches in high-energy and condensed matter physics. Numerous studies have focused on low-dimensional magnetic topological solitons such as one-dimensional (1D) magnetic domain walls¹ and 2D magnetic skyrmions², and explicitly revealed the promising applications of topological solitons in spintronic devices. 3D topological textures with more complex features have also been attracting considerable interest in numerous branches of physics. For example, hopfions, as the most classic 3D topological solitons originally raised by Heinz Hopf in 1931³, have been proved to be of great significance in chiral magnets, quantum fields, cosmology, fluid dynamics and liquid crystals.

Recently, photonic counterparts of topological solitons have aroused intense interest of researchers. Optical skyrmions and optical hopfions have been experimentally demonstrated and considered to have potential applications in metrology, information encoding, and optical communication^{4,5}. However, all the current researches on photonic topological solitons are restricted to spatial dimensions and lose sight of temporal or dynamic information of such structures, and hence manipulating corresponding light-matter interactions spatiotemporally remains challenging.

In the article in *eLight*⁵, the research group led by Prof. Qiwen Zhan from University of Shanghai for Science and Technology theoretically designed and experimentally demonstrated dynamic scalar optical hopfions weaved by nested equiphase lines in the shape of a toroidal vortex. This spatiotemporally varied optical hopfion is proposed through an analytical expression as an approximate solution to Maxwell's equations. Numerical simulations and experimental data demonstrate that the equiphase lines are disjoint and linked closed loops in the form of links and knots with a linking number determined by the Hopf invariant. As shown in Fig. 1, each closed loop painted in one specific color corresponds to a point in the parameter space and to a circle on unit sphere in 4D space. Each closed loop is an equiphase line and all equiphase loops form complete tori that fill up the entire 3D space.

The key to experimentally generate such dynamic scalar optical hopfions is to manipulate poloidal and toroidal spiral phases in toroidal coordinates. In this regard, three spatial light modulators are utilized to generate spatiotemporal optical vortex and flexibly control the winding numbers of the optical hopfions. The authors perform 2D phase measurements of the poloidal phase through interfering the hopfion wave packet with a transform-limited reference pulse split from the source. The

Correspondence: Xiangping Li (xiangpingli@jnu.edu.cn)

¹Guangdong Provincial Key Laboratory of Optical Fiber Sensing and Communications, Institute of Photonics Technology, Jinan University, Guangzhou, China

© The Author(s) 2023



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.






Fig. 1 Schematic of the Hopf fibration and scalar hopfion by spatiotemporal beam shaping. The equiphase lines in a scalar hopfion present the topological features of the Hopf fibration

measured poloidal phase distributed in a spiral pattern in the spatiotemporal domain is well consistent with the theoretical predictions.

As the obtained dynamic scalar optical hopfions manifest as spatiotemporally structured pulses propagating in space-time, they may provide a photonic platform for studying particle-like topological states, and may find promising potential applications in light-matter interactions with spatiotemporal mode excitation, and in optical communications as high-dimensional information carriers.

Published online: 02 March 2023

References

1. Parkin, S. S. P. et al. Magnetic domain-wall racetrack memory. *Science* **320**, 190–194 (2008).
2. Tsesses, S. et al. Optical skyrmion lattice in evanescent electromagnetic fields. *Science* **361**, 993–996 (2018).
3. Hopf, H. Über die Abbildungen der dreidimensionalen Sphäre auf die Kugelfläche. *Math. Ann.* **104**, 637–665, <https://doi.org/10.1007/BF01457962> (1931).
4. Sugic, D. et al. Particle-like topologies in light. *Nat. Commun.* **12**, 6785 (2021).
5. Wan, C. H. et al. Scalar optical hopfions. *eLight* **2**, 22, <https://doi.org/10.1186/s43593-022-00030-2> (2022).